References

1. Abraham D. J.; (2003); Burger's., Medicinal Chemistry & Drug Discovery**;** 6(4); 24-30.
2. Oliver K.; and Alexander S.; (2005); Microwaves in Organic and Medicinal Chemistry; 3; 144-156.
3. Xiao,Fang. Liu; Chang,Ji. Zheng; Liang,Peng. Sun; Xue,Kun. Liu; and Hu,Ri. Piao; (2011) Synthesis of new chalcone derivatives bearing 2,4- thiazolidinedione and benzoic acid moieties as potential anti bacterial agents; European Journal of Medicinal Chemistry**;** 46(8); 3469-3473
4. Bhagat T, M.; Swamy D, K.; Badne S, G. and Kuberkar S,V.; (2011)**,** Synthesis and antibacterial activity of 4- thiazolidinone containing benzothiazolyl moiety; Rasayan journal of chemistry**;** 4; 24-28.
5. Youfu L.; Liang M. and Hao Z.; (2010); Discovery of (z)-5-(4-methoxy benzylidene) thiazolidine-2,4-dione, a readily available and orally active glitazone for the treatment of concansvslin A-induced acute injury of BALB/c mice; Journal of medicinal chemistry; 53; 273-281.
6. Amal M. Youssef; M. Sydney White; and Erika B. Villanueva.; (2010); Synthesis and Biological evaluation of novel pyrazolyl-2,4-thiazolidinediones as anti-inflammatory and neuroprotective agents; Bioorganic & Medicinal Chemistry; 18(5); 2019-2028.
7. Wu Ying; Tai Hsin-Hsiung; and Cho Hoon; (2010); Synthesis and SAR of thiazolidinedione derivatives as 15-PGDH inhibitors.; Bioorganic & Medicinal Chemistry.; 18(5);1428-1433.
8. Carbone Vincenzo; Giglio Marco; Chung Roland; and Trevor Huyton;(2010); Structure of aldehyde reductase in ternary complex with a 5-arylidene-2,4-thiazolidinedione aldose reductase inhibitor; European Journal of Chemistry; 45(3); 1140-1145.
9. Pattan S. R; Suresh C. H; and Pujar V.D; (2009); Synthesis and Antidiabetic activity of 2-amino[5'(4-sulphonylbenzylidene-2,4-thiazolidinedione]-7-chloro-6-flurobenzothiazole; Indian Journal of Chemistry; 44B; 2404-2408.
10. .Borkar M. R; Sachan N.; and Shivajirao S; (2009); Synthesis of N-substituted 2,4-thiazolidinediones and hypoglycemic activity and QSAR Analysis;. Indian journal of Heterocyclic Chemistry, 18295-300.

11. Pattan S.R; and Kekare P.; (2009); Synthesis & Evaluation of some new thiazolidinediones derivatives for their Antidiabetic activities; Asian J.Research Chem; 2(2);256-261.
12. Shelze K.F; Sapkal S.B; and Madje B.R; (2009); Ionic liquid promoted on efficient synthesis of 5-Arylidene-2,4-thiazolidinedione; Bulletin of the Catalysis Society of India; 8; 30-34.
13. Shital Mahalle; Dinesh Ligampalle; and Ramroa Mane; (2009); Microwave-assisted synthesis of some 2,4-thiazolidinediones derivatives; Heteroatom Chemistry; 20(3) ;1695-1700.
14. Armagan Onal; (2009); Spectrophotomeric and HPLC determinations of anti-diabetic drugs, Rosiglitazone maleate and merformin hydrochloride, in pure form and in pharmaceutical preparations; European Journal of Medicinal chemistry; 44; 4998-5005.
15. Murugan R.; Anbazhagan S.; and Sriman S.N.; (2009); Synthesis and invivo antidiabetic activity of novel dispiropyrrolidines through [3+2] cycloaddition rections with thiazolidinedione and rhodanine derivatives; European Journal of Medicinal chemistry; 44; 3271-3279.
16. Sundriyal S.; and Prasad V.B.; (2009); Important pharmacophores features of pan PPAR agonists: common chemical feature analysis and virtual screening; European Journal of Medicinal chemistry; 44; 3488-3495.
17. Mohammed S.A; Gaby E.I; and Gammel A.M; (2009); Synthesis, characterization and in vitro antimicrobial activity of novel 2-thioxo-4-thiazolidinones and 4,4'-bis (2-thioxo-4-thiazolidinone-3-yl)diphenylsulfones; European Journal of Medicinal chemistry; 44; 4148-4152.
18. Liu W.; Liu K.; Wood B.H.; (2009); Discovery of a Peroxisome Proliferator activated receptor γ (PPARγ) modulator with balanced PPARα activity fro the treatment of type 2 diabetes and dyslipidemia; Journal of medicinal chemistry; 52; 4443-4453.
19. Sudarshan K.V.; Manish M.; Kumar H.; and Tripathi K.; (2009); Effect of Pioglitazone and Rosiglitazone on mediators of endothelial dysfunction, markers of angiogenesis and inflammatory cytokines in type 2 diabetes; Acta Diabetol 46; 27-33.

20. Dundar B.O; Mentese A.; and Verspohl V. M; (2008); Synthesis and Antidiabetic activity of some new thiazolyl-2, 4-thiazolidinediones; Arzenim-Forsch/Drug Res*;* 58(3); 131-135.
21. Dundar B.O.; Evranos B.; and Das-Evicmen N.;(2008); Synthesis and Aldose reductase inhibitory activity of some new chromonyl-2,4-thiazolidinediones; European journal of medicinal chemistry; 43; 2412-2417.
22. Pattan S. R.; Khade A. B.; and Buktitagar A.; (2008); Synthesis and evaluation of some new 2,4-thiazolidinediones and their antidiabetic activity; Indian Journal of Heterocyclic Chemistry; 17; 299-300.
23. Hong-Yan Liu; and Shu.Shen Lin; (2008); COMFA and COMSIA analysis of 2,4-thiazolidinediones derivatives as aldol reductase inhibitors; J. Mol. Model; 69-72.
24. Thomas C.; and Emilie D.;(2008); Regulation of thiazolidinediones and implication in glyceroneogenesis in adipose tissue; Diabetes; 57; 68-74.
25. Falck J. R.; and Shuanhu G.; (2008); Elactrophilic α-thiocyanation of chiral and achiral N-acyl carboximides. A convenient route tp 5-substituted and 5,5-disubstituted 2,4-thiazolidinediones; Bioorganic & Medicinal Chemistry; 18; 1760-1771.
26. Jian Yang; and Shuo Wei; (2008); Pharmacological exploitation of the Peroxisome proliferators activated receptor γ agonist Ciglitazone to develop a novel class of androgen receptor-Ablative agent; Journal of Medicinal Chemistry; 51; 2100-2107.
27. Pattan S.R; Alagwadi K.R; Bhat A.R; and Reddy V.V.K; (2007); synthesis and evaluation of aome 5-[1[(4-(4-substituted –phenylamino)-met-(z)-ylidene]-Thiazolidine-2,4-diones for antitubercular activity; Indian drugs; 45(7); 256-262.
28. Ameya A.C.; and Pai N.R.; (2007); Synthesis & antihyperglycemic activity of 2,4-thiazolidinediones; Indian Journal of heterocyclic activity; 17; 45-48.
29. Sachan N.; and Shivajirao S. K; (2007); Synthesis, antihyperglycemic activity and QSAR of 5-Benzylidene-2, 4-thiazolidinediones; Indian Journal of Heterocyclic Chemistry; 17; 57-62.
30. Pattan S.R.; Reddy V.V.K.; Pawar P.D.; and Khade A.B.; (2007); Synthesis and anti-inflammatory evaluation of (4-phenylamino benzylidene (thiazolidine-2,4-dione)derivatives; Indian Drugs; 44(4); 143-147.

31. Ali M.A.; Gamal E. S.; and Mahfouz N. M.; (2007); Synthesis and three dimensional Qualitative Structure Selectivity relationship of 3,5-disubstituted-2,4-thiazolidinedione derivatives as COX2 inhibitors; Arch. Pharm. Res.; 30(10); 1186-1204.
32. Dundar B.O.; Unlusoy C.M.; and Verspohl E. J.; (2007); Synthesis and Antidiabetic activity of some new Chromonyl-2,4-thiazolidinediones; Arzenim-Forsch/Drug Res.; 50(II); 626-630.
33. Shiradkar R. M.; and Ghodake M.; (2007); Synthesis and anticonvulsant activity of clubbed Thiazolidinone-barbituric acid and Thiazolidinone-triazole derivatives; ARKIVOC; (xiv); 58-74.
34. Wei S.; and Lin Li Fang; (2007); Thiazolidinediones modulate the expression of β-catenin & other cell cycle regulatory proteins by targeting the F-box proteins of SKp-1-Cul1-F-box protein E 3 ubliquitin Ligase independently of Peroxisome proliferators-activated receptor γ; Molecular pharmacology; 72; 725-733.

`